



Resilience in a Changing Climate: Advancing Research on Groundwater for Equity

Vietnamese Mekong Delta (VMD)

Groundwater Profile

February 2025



76% - 83% of households surveyed by the RECHARGE team relied on groundwater for drinking and domestic purposes (in wet and dry seasons respectively).



50% Of VMD households are dependent on groundwater for their livelihoods.

Drinking water use

In rural Vietnam, bottled water is the primary drinking water source, followed by rainwater.





Fig. 1 – % Volume of VMD water use by sector (Ha et. al. 2015).

Fig. 2 – Primary drinking water sources in the VMD (Viet Nam MICS, 2021).

Drinking water use by ethnicity

Khmer communities are more reliant on private supplies such as rainwater, wells and boreholes




Fig. 3 – Primary dry season drinking water source by ethnicity (UTS-ISF, 2024).

* Bottled water and piped water can be sourced from groundwater.

Drinking water use by wealth quintile

Higher-income households have a higher rate of access to piped supplies and lower income households rely more on rainwater and private groundwater.




Fig. 3 – Use of drinking water sources in VMD by wealth quintiles (Viet Nam MICS, 2021).




Fig. 4 – Spatially interpolated recent (1996–2017) trends of GWLs of the seven different aquifers in the VMD (Duy et al. 2021).

Groundwater level changes

- Potable water aquifers in the VMD are decreasing at a rate of 85cm per year (An Tran et al. 2022).
- Groundwater extraction exceeds groundwater recharge in these aquifers by 35% (Van et al., 2023).

Groundwater salinity changes

- Declining water levels in potable aquifers are leading to increased saline intrusion from shallow aquifers.
- Saline intrusion is exacerbated by climate change-induced sea level rise.
- High chloride levels are predicted in the main potable aquifer to a depth of 150m by 2030, exceeding the Viet Nam technical regulation on drinking water quality of 0.25g/L (An Tran et al., 2022).




Fig. 5 – Cross-section of the seven groundwater aquifers in the VMD (Doan et al., 2016 as cited by Gunnik et al., 2021).

Water reliability

Bottled water and rainwater are considered by households to be the most reliable drinking water sources with significantly lower perceived reliability for piped groundwater and private groundwater sources.




Fig. 6 – Percentage of households reporting water as always available by source (UITS-ISF, 2024).

Water quality

Water quality is a concern across most sources. When tested for *E. coli*, boreholes and protected wells had the lowest contamination rate among at 13%, lower than bottled water (26%). Surface water had the highest levels of *E. coli* at 100%.




Fig. 7 – Proportion of sampled drinking water sources testing positive for *E. coli* colony forming units (Viet Nam MICS, 2021).



Borehole registration

Only 13% of private boreholes are registered with government authorities limiting monitoring and planning of extraction.




Fig. 8 – Percentage of private boreholes users reporting registering the borehole with local authorities (UTS-ISF, 2024)



Challenges

- Groundwater levels are decreasing across the VMD as demand exceeds recharge rates
- Salinity is increasing across VMD groundwater aquifers further reducing the availability of acceptable drinking water
- *E. coli* contamination is significant, and household treatment is required to ensure safety
- Lower wealth households and Khmer households have significantly lower levels of access to on-premises piped water supplies
- Low levels of borehole registration are limiting monitoring of extraction levels

References

- An Tran, D., Tsujimura, M., Pham, H. V., Nguyen, T. V., Huu Ho, L., Le Vo, P., Quang Ha, K., Duc Dang, T., Van Binh, D., & Doan, Q.-V. (2022). Intensified salinity intrusion in coastal aquifers due to groundwater overextraction: a case study in the Mekong Delta, Vietnam. *Environmental Science and Pollution Research*.
- Gunnink, J. L., Pham, H. V., Oude Essink, G. H. P., and Bierkens, M. F. P. (2021) The three-dimensional groundwater salinity distribution and fresh groundwater volumes in the Mekong Delta, Vietnam, inferred from geostatistical analyses, *Earth Syst. Sci. Data*, 13, 3297–3319
- Ha K, Ngoc NTM, Lee E, Jayakumar R, (2015), Current status and issues of groundwater in the Mekong river basin, Korea Institute of Geoscience and Mineral Resources (KIGAM), CCOP Technical Secretariat and UNESCO Bangkok
- Duy NL, Nguyen TVK, Nguyen DV, Tran AT, Nguyen HT, Heidbüchel I, Merz B, Apel H, (2021), Groundwater dynamics in the Vietnamese Mekong Delta: Trends, memory effects, and response times, *Journal of Hydrology: Regional Studies*, Volume 33, 2021,
- UTS-ISF and CTU. (2024). RECHARGE household survey data.
- Van, T. D., Zhou, Y., Stigter, T. Y., Van, T. P., Hong, H. D., Uyen, T. D., & Tran, V. B. (2023). Sustainable groundwater development in the coastal Tra Vinh province in Vietnam under saltwater intrusion and climate change. *Hydrogeology Journal*,
- Viet Nam MICS (2021). Viet Nam General Statistics Office and UNICEF. 2021. Viet Nam Multiple Indicator Cluster Survey 2021, Survey Findings Report. Ha Noi Viet Nam

Suggested citation

UTS-ISF. *Vietnamese Mekong Delta Groundwater Profile*. University of Technology Sydney – Institute for Sustainable Futures, Sydney, 2025.

All photo credits: Diana Gonzalez (UTS-ISF)

For more information contact naomi.carrard@uts.edu.au